

A novel phase in $SU(3)$ gauge theory with many light fermions

Anna Hasenfratz
University of Colorado



New Horizons for Lattice Computations Workshop
BNL
May 16 2012

In collaboration with A. Cheng, D. Schaich and G. Petropoulos
ArXiv:1111:2317

The many faces of gauge-fermion systems

Renormalization group β function at 2 loops


$$\beta(g^2) = \frac{dg^2}{d \log(\mu^2)} = \frac{b_1}{16\pi^2} g^4 + \frac{b_2}{(16\pi^2)^2} g^6 + \dots$$
$$b_1 = -11 + \frac{2}{3} N_f, \quad b_2 = -102 + \frac{38}{3} N_f$$



The many faces of gauge-fermion systems

Renormalization group β function at 2 loops

$$\beta(g^2) = \frac{dg^2}{d \log(\mu^2)} = \frac{b_1}{16\pi^2} g^4 + \frac{b_2}{(16\pi^2)^2} g^6 + \dots$$
$$b_1 = -11 + \frac{2}{3} N_f, \quad b_2 = -102 + \frac{38}{3} N_f$$



Chirally broken & confining

- QCD with 2+1 (+1) flavors
- Original technicolor candidates


The many faces of gauge-fermion systems

Renormalization group β function at 2 loops

$$\beta(g^2) = \frac{dg^2}{d \log(\mu^2)} = \frac{b_1}{16\pi^2} g^4 + \frac{b_2}{(16\pi^2)^2} g^6 + \dots$$
$$b_1 = -11 + \frac{2}{3} N_f, \quad b_2 = -102 + \frac{38}{3} N_f$$

Conformal

- IRFP where g_0 is irrelevant
- universal anomalous dimension



$$8 < N_f < 16.5$$




The many faces of gauge-fermion systems

Renormalization group β function at 2 loops

$$\beta(g^2) = \frac{dg^2}{d \log(\mu^2)} = \frac{b_1}{16\pi^2} g^4 + \frac{b_2}{(16\pi^2)^2} g^6 + \dots$$
$$b_1 = -11 + \frac{2}{3} N_f,$$
$$b_2 = -102 + \frac{38}{3} N_f$$

What we really want is walking




The gauge coupling changes slowly and the anomalous mass dimension remains large across an extended energy scale



Roadmap for the conformal window

S-D type calculations




- Which models are conformal?
- Which – if any - is walking?
- What is the anomalous mass ?
- S parameter?



Roadmap for the conformal window

S-D type calculations



- What is the best approach to study these systems?
- Do we understand the systematics?

Don't be impatient : QCD has a 25 year head start



A phase with novel symmetry breaking pattern

Strongly coupled systems can lead to unexpected phenomena:

$N_f=12$ and 8 flavors SU(3) with staggered fermions

- 8 flavors: most likely chirally broken
- 12 flavors: still controversial


Goal: study the phase diagram both at **zero and at finite temperature** and contrast the two systems

(Lattice action: it's a good one)




Why Finite temperature ?

QCD like



$$\beta_c \rightarrow \infty \text{ as } N_T \rightarrow \infty$$

Conformal





$$\beta_c \rightarrow \beta_{bulk} \text{ as } N_T \rightarrow \infty$$



The scaling of β_c is a good test of conformality
(possibly)



The phase structure of $N_f=12$




The phase structure of $N_f=12$



Similar transitions were observed previously by INFN/Groeningen group and LHC collaboration




Phase diagram β - m plane $N_f=12$



Intermediate phase bordered by bulk 1st order transitions



Phase diagram β - m plane $N_f=12$



Intermediate phase bordered by bulk 1st order transitions





New symmetry breaking pattern in the IM phase

Single-site shift symmetry (S^4) of the staggered action

$$\begin{aligned}\chi(n) &\rightarrow \xi_\mu(n)\chi(n+\mu), & \xi_\mu(n) &= (-1)^{\sum_{\nu>\mu} n_\nu} \\ U_\mu(n) &\rightarrow U_\mu(n+\mu),\end{aligned}$$

is broken \rightarrow plaquette expectation value is “striped”




New symmetry breaking pattern

Order parameter I:

$$\Delta P_\mu = \langle \text{Re Tr } \square_n - \text{Re Tr } \square_{n+\mu} \rangle_{n_\mu \text{ even}}$$

Plaquette on even & odd time slices are different – this is on the background gauge configuration!



$\beta = 2.6$ IM phase

$\beta = 2.7$ weak coupling phase





New symmetry breaking pattern

S^4 breaking occurs at the fermionic level:

Order parameter –II: link difference

$$\Delta L_\mu = \langle \alpha_\mu(n) \bar{\chi}(n) U_\mu(n) \chi(n + \mu) - \alpha_\mu(n + \mu) \bar{\chi}(n + \mu) U_\mu(n + \mu) \chi(n + 2\mu) \rangle_{n_\mu \text{ ev}}$$



β=2.7 weak phase



New symmetry breaking pattern


- Single-site shift symmetry is exact in the action.
- Both ΔP and ΔL are order parameters of S^4
- When S^4 is broken, the phase has to be separated by a “real” phase transition
- The S^4 broken (~~S^4~~) phase cannot go away with the volume
- S^4 is related to taste so this could be a special taste breaking

What are the physical properties of the S^4_b phase?



The Polyakov line

Is it deconfining? Polyakov line is very noisy but the **blocked Poly line** is sensitive:




Blocked Poly line is measured on RG blocked lattices:

- improved Poly line
- or
- Poly line on renormalized trajectory, after removing UV fluctuations

The static potential \rightarrow Confinement!

Static potential on $12^3, 16^3$ volumes (no volume dependence!) shows a linear term: $r_0 = 2.1 - 2.7$, $\sqrt{\sigma} = 0.40 - 0.48$



$\beta = 2.6$ – IM phase


$\beta = 2.7$ – weak coupling

No comment about the weak coupling side from the potential – too small volume



Chiral properties : Dirac eigenvalue spectrum

4 different volumes, 12 eigenvalues each:



Qualitatively different – quantitative description?



Dirac eigenvalue spectrum

- RMT predictions require knowledge of the dynamics
- Simple volume scaling is more general. In the chiral limit

$$\rho(\lambda) \sim (\lambda - \lambda_0)^\alpha \quad \lambda_0 : \text{soft edge}$$

$$\int_m^n \rho(\lambda) d\lambda = \frac{n - m}{V} + O(1/V^2)$$

$$\lambda_n - \lambda_0 \sim \left(\frac{n - x_0}{V} \right)^{1/(\alpha+1)}, \quad \frac{D}{\alpha+1} = y_m$$

- Conformal IRFP : $y_m = 1 + \gamma^*$, : independent of V , β
- Confining system : $y_m = 1 + \gamma(L)$: in volumes smaller than the confinement scale





Dirac eigenvalue spectrum


Fit in S^4b phase ($\beta=2.6$):

Needs soft edge $\lambda_0=0.0175$ with $\alpha=0.6(1)$ (RMT prediction: $\alpha=1/2$)

Very chiral symmetric! (Even $U(1)_A$ restoring)



Scaling of Dirac eigenmodes, $N_f = 12$, weak coupling side



Two coupling values, four volumes with 30 eigenmodes each are consistent, predict $y_m = 0.47(2)$

Very efficient method to determine the anomalous dimension (we need larger volumes, other β , more eigenmodes to reduce error)



The S^4_b phase


Is most likely a lattice artifact

- Breaks single-site translational symmetry
- It is confining with small correlation length
- It is chirally symmetric
- (yes, it would violate the 't Hooft anomaly matching in the continuum)

What happens at finite temperature?



Phase structure at finite T , $N_f=12$



$N_T=12$, 16 transitions

- indistinguishable
- run into the S^4b phase


The weak coupling, $m < 0.03$ region is either

- conformal or
- has $N_T > 16$

Preliminary; more data in progress



Improved Polyakov line



Poly line is difficult to measure at large N_t as it decays exponentially

$$\langle P \rangle \sim \exp(-f \cdot N_t)$$

Smeared (HYP) or blocked Poly gives a better signal



S⁴b phase in other systems


There are signs of the intermediate phase with other $N_f=12$ actions

Is it unique to $N_f=12$?

No.



Phase structure with $N_f=8$



The S⁴b phase, bordered by bulk transitions, is present
→ S⁴b does not imply conformality

But $N_T=12, 16$ transitions behave differently: move toward $g^2 = 0$ as N_T increases

This is consistent with confinement

Preliminary; more data in progress



Summary & Outlook

These systems are complicated and have strange (strong coupling) lattice artifacts

Progress is steady but we need better understanding

The S^4_b phase is present with 2 sets of staggered fermions. Could it show up in 2+1 flavor simulations?

